




Immunization Recommendations for Children Aged 0 Through 18 Years

KEY

HepB = to be performed  = range of recommended ages to be performed  = certain high-risk groups  = catch-up immunization

AGE ▶	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years	7-10 years	11-12 years	13-18 years
▼ VACCINE														
Hepatitis B	HepB	HepB			HepB							HepB Series		
Rotavirus			RV	RV	RV									
Tetanus, Diphtheria, Pertussis (Td/Tdap)			TDaP	TDaP	TDaP		TDaP				TDaP		TDaP	TDaP
<i>Haemophilus influenzae</i> type b			Hib	Hib	Hib	Hib								
Pneumococcal			PCV	PCV	PCV	PCV				PPSV				
Inactivated Poliovirus			IPV	IPV	IPV						IPV	IPV		
Influenza					Influenza (Yearly)									
Measles, Mumps, Rubella (MMR)						MMR					MMR	MMR series		
Varicella (Chicken Pox)						Varicella					Varicella	Varicella series		
Hepatitis A						HepA (2 doses)				HepA series				
Meningococcal										MCV		MCV	MCV	
Human Papillomavirus												see footnote 3	HPV (3 doses)	HPV series

For complete statements, refer to the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/recs/acip), the American Academy of Pediatrics (<http://www.aap.org>), and the American Academy of Family Physicians (<http://www.aafp.org>).

The Ohio State University Health Plan Inc. created this immunization schedule based on the Recommended Immunization Schedule for Persons Aged 0 Through 6 Years, United States, 2009, and Recommended Immunization Schedule for Persons Aged 7 Through 18 Years, United States, 2009, published in the Morbidity and Mortality Weekly Report on January 9, 2009 (MMWR 2008;57[53]).